

**Table 3.1.1: Additional\* vaccines recommended for Indigenous persons, due to their higher risk of disease**

Vaccine	Recommendation for Indigenous persons
BCG	Neonates living in areas of high TB incidence <sup>†</sup> 1 dose
Hepatitis A	Children resident in the Northern Territory, Queensland, South Australia and Western Australia 2 doses in the 2nd year of life <sup>‡</sup>
Hepatitis B	Adults who have not previously been vaccinated against hepatitis B and are non-immune
Influenza	All persons aged $\geq 6$ months <sup>§</sup> Annual vaccination
Pneumococcal conjugate (13vPCV)	Children resident in the Northern Territory, Queensland, South Australia and Western Australia Booster dose in 2nd year of life in addition to primary course <sup>‡</sup>
Pneumococcal polysaccharide (23vPPV)	Persons aged 15–49 years with underlying conditions increasing the risk of IPD <sup>¶</sup> All persons aged $\geq 50$ years <sup>¶</sup>

\* In addition to those vaccines recommended for all Australians or those in particular medical, occupational, behavioural or other risk groups.

† Northern Territory, Queensland, northern South Australia

‡ Exact ages may differ between jurisdictions.

§ Refer to 4.7 *Influenza*.

¶ Refer to 4.13 *Pneumococcal disease* for recommendations on revaccination.